



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A21D 2/02, A23K 1/18	A1	(11) International Publication Number: WO 00/30456 (43) International Publication Date: 2 June 2000 (02.06.00)
<p>(21) International Application Number: PCT/US99/26640</p> <p>(22) International Filing Date: 12 November 1999 (12.11.99)</p> <p>(30) Priority Data: 60/109,686 24 November 1998 (24.11.98) US 09/428,631 27 October 1999 (27.10.99) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/428,631 (CIP) Filed on 27 October 1999 (27.10.99)</p> <p>(71) Applicant (for all designated States except US): RALSTON PURINA COMPANY [US/US]; Checkerboard Square, St. Louis, MO 63164 (US).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): BRUNNER, John, J. [US/US]; 1314 Briarcreek, Kirkwood, MO 63122 (US).</p> <p>(74) Agents: BEULICK, John, S. et al.; Armstrong Teasdale LLP, Suite 2600, One Metropolitan Square, St. Louis, MO 63102-2740 (US).</p>	<p>(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU (Petty patent), AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>	
(54) Title: METHODS AND COMPOSITIONS FOR ENHANCING PALATABILITY OF PET FOOD		
<p>(57) Abstract</p> <p>Compositions and methods for enhancing the palatability of dry and semi-dry pet food compositions, particularly extruded cat food compositions, are described. In one embodiment, the palatability enhancer is a dry cat food coating including tetrasodium pyrophosphate at about 0.1 % to about 1.0 % by weight of the finished pet food product. The palatability enhancer is applied after the extrusion process to the extruded particles or pieces of pet food.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS AND COMPOSITIONS FOR ENHANCING PALATABILITY OF PET FOOD

Cross Reference to Related Application

This application claims the benefit of U.S. Provisional Application No.
5 60/109,686, filed 11/24/98.

Field of the Invention

This invention relates generally to methods and compositions for enhancing the palatability of pet foods, and more particularly to compositions and methods for enhancing the palatability of dry and semi-dry cat food.

10

Background of the Invention

Pet food manufacturers have a long-standing desire to provide foods which combine high nutritional value, and resistance to decomposition and bacterial contamination, with low production costs. In addition, and with
15 particular regard to cat foods, pet food manufacturers desire a high degree of palatability which can be attained at low cost.

The three main types of pet foods embody the desired characteristics to varying degrees. Canned or high-moisture (greater than 50% moisture) foods are typically all meat and therefore generally highly palatable to animals. However,
20 such foods are typically costly to produce and package, require the use of preservatives to inhibit decomposition and contamination, and are relatively less nutritious per unit weight than other pet food products. Dry or low-moisture content pet foods (less than 15% moisture) are generally highly nutritious, may be inexpensively packaged, and are highly convenient to store and use.
25 However, dry or low-moisture foods are generally the least palatable of pet foods. Semi-dry or intermediate moisture content pet foods (about 15% to about 50% moisture) are generally more palatable than dry foods, have higher nutritional value, and are more inexpensively packaged and more conveniently

- 2 -

used than high moisture foods. However, intermediate moisture content foods may be more susceptible to bacterial contamination than dry pet foods, and are generally less palatable than high moisture content foods.

Dry and semi-dry foods are generally preferred by manufacturers and pet owners alike because such foods are more convenient to store and use, and because of the higher nutritional content than high-moisture pet foods. However, many animals, and particularly cats, are picky eaters desiring a high degree of palatability. There remains a need for a low-moisture pet food with a high degree of palatability.

Known palatability enhancers for low moisture pet foods include various surface coating compositions. For example, known palatability enhancers include surface coatings of phosphoric acid; a combination of fat with hexamic, phosphoric or citric acid; and a mixture of phosphoric acid and citric acid. Palatability may also be enhanced by the application of liquid or dry flavorings as a coating. Liquid or dry animal digest coatings are widely used to enhance palatability, and animal digest is also used in a liquid combination with phosphoric acid. Dry animal digest may be used in a two step coating process with liquid phosphoric acid. Dry application of sodium acid phosphate, which may be used in combination with animal digest, is also known.

However, the known compositions and methods remain limited in the resulting degree of palatability. It would be desirable to provide a composition and method for further enhancing the palatability of dry and semi-dry pet foods. It would also be desirable to provide such a composition with a high degree of palatability for cats. It would be further desirable to provide such a composition and method which enhances dry and semi-dry pet food palatability at a relatively low production cost. It would be still further desirable to provide such a palatability enhancer which is robust among different formulations.

- 3 -

Summary of the Invention

These and other objects may be attained by a pet food palatability enhancer which includes tetrasodium pyrophosphate. The palatability enhancer, and methods for enhancing the palatability of pet foods, generally relate to dry and semi-dry pet foods including a nutritionally balanced mixture of proteinaceous and farinaceous materials. Generally, the palatability enhancer is a dry formulation including from about 0.1% up to about 99% by weight tetrasodium pyrophosphate. In one embodiment, the palatability enhancer is formulated by adding tetrasodium pyrophosphate in dry powdered form to a dry composition for application to a dry pet food. The dry composition typically also includes readily commercially available ingredients such as dried whey, dried animal digest, and yeast. The amount of tetrasodium pyrophosphate added to the dry composition is adjusted so that tetrasodium pyrophosphate contributes about 0.1% to about 1.0% by weight of the finished dry pet food.

In use and in one embodiment, the palatability enhancer is applied to particles or pieces of extruded, dry or semi-dry cat food. More specifically, the particles or pieces are added to a coating drum and molten animal fat and other liquids are sprayed over the dry pieces. The palatability enhancer is applied to the pieces and the pieces are tumbled in the drum to thoroughly coat the pieces.

The palatability enhancer improves the palatability of dry and semi-dry pet foods. The palatability enhancer is particularly effective for enhancing the palatability to cats of dry and semi-dry cat foods. Further, the palatability enhancer may be added to commercially available dry and semi-dry cat foods or other pet and animal foods at a relatively low production cost.

25

Detailed Description

The palatability enhancer generally is a dry composition including from about 0.1% up to about 99% by weight tetrasodium pyrophosphate. The

- 4 -

palatability enhancer is generally externally applied, using dry application methods, to commercially available, dry or semi-dry pet foods.

More specifically, the palatability enhancer and methods for enhancing the palatability of dry and semi-dry pet foods, generally relate to a nutritionally
5 balanced mixture of proteinaceous and farinaceous materials having a moisture content of about 50% or less by weight. Such mixtures are known as dry or semi-dry pet foods to those skilled in the art, and the palatability enhancer is applied to pieces of the dry or semi-dry food. The pet food compositions described herein are not intended to be limited to a specific listing of ingredients
10 because such ingredients will depend on such factors as, for example, the desired nutritional balance for the specific type of pet, and availability of ingredients to the manufacturer. In addition to the proteinaceous and farinaceous materials, the pet food composition generally may include vitamins, minerals, and other additives such as flavorings, preservatives, emulsifiers and humectants. The
15 nutritional balance, including the relative proportions of vitamins, minerals, fat, protein and carbohydrate, is determined according to dietary standards known in the veterinary art. For example, the nutritional balance of a cat food composition is determined according to the known dietary requirements for cats.

The proteinaceous material may include any material having a protein
20 content of at least about 15% by weight including vegetable proteins such as soybean, cotton seed, and peanut; animal proteins such as casein, albumin, and fresh animal tissue including fresh meat tissue and fresh fish tissue; and dried or rendered meals such as fish meal, poultry meal, meat meal, bone meal and the like. Other types of suitable proteinaceous materials include wheat gluten or corn
25 gluten, and microbial proteins such as yeast.

The farinaceous material may be defined as any material having a protein content of less than about 15% by weight and containing a substantial proportion of starches or carbohydrates, including grains such as corn, milo, alfalfa, wheat, barley, rice, soy hulls, and other grains having low protein content. In addition

- 5 -

to the proteinaceous and farinaceous materials, other materials such as whey and other dairy by-products, as well as other carbohydrates may be added. In addition, known flavorings including, for example, corn syrup or molasses, may be added.

5 In one exemplary embodiment, the palatability enhancer is applied to a dry cat food composition. Generally, the term cat food composition as used herein applies to commercially sold, nutritionally balanced food compositions which are intended to provide substantially the sole diet for a cat. Thus, such compositions may be described as having a minimum protein content at which cat
10 health is maintained. However, the minimum protein content of the food varies according to the age and breeding status for the animal. For example, a nutritionally balanced cat food composition for breeding females and kittens requires a minimum protein content of at least about 28% by weight on a dry matter basis. A nutritionally balanced cat food composition for non-breeding and
15 adult cats requires a minimum protein content of about 26% by weight on a dry matter basis. More typically, the protein content of commercially available cat food compositions for adult, non-breeding cats is about 30% by weight on a dry matter basis, to insure that the food meets the nutritional requirements of any cat.

For example, a typical formula well known in the art for a dry cat food
20 composition to which the palatability enhancer is applied is as follows (in percent by weight): about 0 - 70% grain-based meal or flour, such as corn, wheat, barley or rice; about 0 - 30% animal by-product meal, such as poultry or beef meal; about 0 - 25% corn gluten meal; about 0 - 25% fresh animal tissue, such as poultry or beef tissue; about 0 - 25% soybean meal or flour; about 0 - 10%
25 animal fat; about 0 - 20% seafood-based meal; about 0 - 25% fresh fish tissue; about 0 - 10% high fructose corn syrup; about 0 - 10% dried molasses; about 0 - 1.5% phosphoric acid; and about 0 - 1.5% citric acid. Vitamins and minerals are added according to known American Association of Feed Control Officials (AAFCO) Cat Food Nutrient profiles, and include calcium carbonate, potassium

- 6 -

chloride, sodium chloride, choline chloride, taurine, zinc oxide, ferrous sulfate, vitamin E, vitamin A, vitamin B₁₂, vitamin D₃, riboflavin, niacin, calcium pantothenate, biotin, thiamine mononitrate, copper sulfate, folic acid, pyroxidine hydrochloride, calcium iodate, and menadione sodium bisulfite complex (a source
5 of vitamin K activity).

In addition, the palatability enhancer relates to other pet foods and animal foods in general without regard to the protein content which typically varies according to species, breeding status, and age, among other factors. For example, the palatability enhancer may be used with a dry or semi-dry dog food
10 composition for non-breeding, adult dogs, which requires a minimum protein content of about 18% by weight on a dry matter basis. Similarly, the palatability enhancer may be applied to a dry or semi-dry puppy food having a minimum protein content of about 22% by weight on a dry matter basis. The palatability enhancer may also be used with other dry and semi-dry foods of varying protein
15 content, and with foods for other animals such as livestock and research animals.

The palatability enhancer generally includes tetrasodium pyrophosphate and is formulated as a dry mixture in powdered, granulated or encapsulated form. Tetrasodium pyrophosphate is available in dry, powdered form from, for example, Solutia of St. Louis, Missouri, and is combined with other dry
20 ingredients including known palatability enhancing ingredients and preservatives. For example, other suitable ingredients for the palatability enhancer include various nucleotides such as inosine monophosphate (IMP), guanosine triphosphate (GTP) and adenosine triphosphate (ATP); amino acids such as alanine and lysine; other inorganic salts such as NaCl, monosodium phosphate, and dicalcium
25 phosphate; and organic materials such as whey, yeast and animal by-products digest. The palatability enhancer includes about 0.1% up to about 99% by weight tetrasodium pyrophosphate. In one embodiment, the amount of tetrasodium pyrophosphate is adjusted so that tetrasodium pyrophosphate contributes about 0.1% to about 1.0% by weight to the finished pet food,

- 7 -

depending on the relative amount of the palatability enhancer added to the finished pet food. Generally, the palatability enhancer contributes from about 0.5% to about 3.0% by weight to the finished cat food product. However, both the exact proportion of tetrasodium pyrophosphate in the palatability enhancer, as well as the relative proportion of the palatability enhancer to the finished cat food product, may be varied and are limited by such factors as relative availability of the ingredients, cost, and health of the animal. In addition, the palatability enhancer described herein is not intended to be limited to a specific listing of ingredients because such ingredients will similarly depend on availability and cost of ingredients to the manufacturer.

To make a dry formulation of the palatability enhancer, for example, commercially available dry ingredients, including tetrasodium pyrophosphate, various nucleotides, amino acids, inorganic salts and organic materials are combined in the desired proportions in a batch mixer and blended to homogeneity. In one embodiment, a dry formulation of the palatability enhancer typically includes about 15 - 25 weight % tetrasodium pyrophosphate, about 0.1 - 3.0 weight % amino acid, about 35 - 50 weight % animal by-products such as dried animal digest, and about 35 - 50 weight % microbial protein such as brewer's yeast. For example, in one embodiment the palatability enhancer includes about 18 weight % tetrasodium pyrophosphate or, in another embodiment, about 25 weight % tetrasodium pyrophosphate. In one specific exemplary embodiment, the palatability enhancer includes about 39.7 weight % dried brewer's yeast, about 39.7 weight % dried chicken liver, about 19.8% tetrasodium pyrophosphate, and about 0.8 weight % L-alanine.

To make a liquid formulation of the palatability enhancer, for example, commercially available liquid ingredients are combined in a mixer. Wet ingredients such as animal tissue are ground or emulsified to a slurry and the liquid ingredients are combined with the slurry. A commercially available protease is added to the slurry to hydrolyze the proteins, and later inactivated

- 8 -

with heat, acid or another method. Preservatives such as sorbates are also added to the slurry. Water is added to adjust the viscosity and the solids content of the slurry to between about 10% and 50% to facilitate the application process as described below. The tetrasodium pyrophosphate, and other dry ingredients as
5 desired, are added to the slurry prior to application. A liquid formulation of the palatability enhancer includes about 5 - 45 weight % tetrasodium pyrophosphate; about 1 - 50 weight % animal tissue such as beef hearts or livers, or chicken hearts or livers; about 0.1 - 3.0 weight % amino acid such as alanine or lysine; about 0.1 - 1.0 weight % preservatives such as sorbic acid and about 0.1 - 2.0
10 weight % enzymes such as protease. For example, one embodiment of the liquid formulation includes about 14 - 20 weight % tetrasodium pyrophosphate.

In use, the palatability enhancer is applied to the particles or pieces of dry or semi-dry pet food such as, for example, dry extruded kibbles as known in the art. The palatability enhancer is applied, for example, by spraying or dusting
15 onto the particles or pieces of the food. More specifically, to make a cat food composition to which the palatability enhancer is applied, proteinaceous and farinaceous materials and additional desired materials as described above are combined to form an admixture and are well blended. The admixture is then transferred to a steam conditioner and subjected to steam and moisture to adjust
20 the moisture content of the admixture to between about 20% and 40% by weight. The conditioned admixture is then cooked under conditions of elevated temperature and pressure in an extruder such as a single screw extruder. The extruder may include a die having a particular shape, such as a fish, cross or circle. The product is segmented into discrete particles or pieces by a rotating
25 cutting knife as the product is extruded.

The particles or pieces are conveyed to a forced air drying system which raises the temperature of the pieces to about 140°F, and reduces the moisture level to about 8% by weight. The dried particles or pieces are then transferred by bulk conveyor to a coating drum and sprayed with animal fat. Other liquids

- 9 -

such as, for example, citric acid or phosphoric acid may alternatively be applied to the pieces, or applied in addition to the animal fat. The pieces are then sprayed or dusted with the palatability enhancer, and tumbled to thoroughly coat the pieces with the palatability enhancer. The pieces are then cooled to ambient
5 temperature and packaged.

The palatability enhancer with tetrasodium pyrophosphate, in both liquid and dry formulations, was tested against known dry cat food coatings. In both formulations, the palatability enhancer was shown to improve palatability over that obtained with known palatability enhancers.

10 In four separate tests described in Examples 1 through 4, eighteen adult cats were each offered a control ration R1, and a test ration R2 in a two bowl test over a period of two days. R1 consisted of dry cat food kibbles of Meow Mix® basal formula, commercially available from Ralston Purina Company, Checkerboard Square, St. Louis, Missouri, 63164, with 2.25% by weight of a
15 typical dry cat coating containing about 0.1 - 3.0 weight % amino acid, about 35 - 50 weight % animal by-products such as dried animal digest, and about 35 - 50 weight % brewer's yeast. R2 was Meow Mix® basal formula kibbles coated with 2.5% by weight of a dry cat coating which also included 0.5% by weight tetrasodium pyrophosphate.

20 Example 1

Of all food consumed over the two days, R2 represented 84.6% of the total, while R1 represented 15.4%, and the ratio of the amount of R2 consumed to the amount of R1 consumed was 5.5. The animals ate 55.3 grams/day/cat of R2, compared with 11.0 grams/day/cat of R1. All eighteen cats preferred R2
25 over R1.

Example 2

Of all food consumed over the two days, R2 represented 82.7% of the

- 10 -

total, while R1 represented 17.3%, and the ratio of the amount of R2 consumed to the amount of R1 consumed was 4.8. The animals ate 58.0 grams/day/cat of R2, compared with 12.0 grams/day/cat of R1. Seventeen cats preferred R2 over R1, while one cat showed no preference.

5

Example 3

Of all food consumed over the two days, R2 represented 85.4% of the total, while R1 represented 14.6%, and the ratio of the amount of R2 consumed to the amount of R1 consumed was 5.9. The animals ate 54.4 grams/day/cat of R2, compared with 8.7 grams/day/cat of R1. Sixteen cats preferred R2 over R1, while one cat showed no preference, and one cat was eliminated for insignificant consumption.

10

Example 4

Of all food consumed over the two days, R2 represented 77.6% of the total, while R1 represented 22.4%, and the ratio of the amount of R2 consumed to the amount of R1 consumed was 3.5. The animals ate 49.2 grams/day/cat of R2, compared with 14.7 grams/day/cat of R1. Sixteen cats preferred R2 over R1, while one cat preferred R1 over R2 and one cat showed no preference.

15

Table 1 summarizes the results of Examples 1 through 4.

TABLE 1

20

Ex.	Ra- tion	most consumed / least consumed	% of all food consumed	Grams eaten /day/cat	# of animals with pref.	# of animals without pref.
1	R1	5.5	15.4	11.0	0	0
	R2		84.6	55.3	18	
2	R1	4.8	17.3	12.0	0	1

- 11 -

	R2		82.7	58.0	17	
3	R1	5.9	14.6	8.7	0	1
	R2		85.4	54.4	16	
4	R1	3.5	22.4	14.7	1	1
	R2		77.6	49.2	16	

In two separate tests set forth in Examples 5 and 6, a control ration coated with liquid animal digest was tested against a test ration coated with liquid animal digest including an amount of tetrasodium pyrophosphate.

Example 5

Eighteen adult cats were each offered a control ration R1, and a test ration R3 in a two bowl test over a period of two days. R1 consisted of Meow Mix® basal formula kibbles coated with liquid animal digest at 3% by weight. R3 was Meow Mix® basal formula kibbles coated with liquid animal digest at 3% by weight, plus tetrasodium pyrophosphate at 0.75% by weight. Of all food consumed over the two days, R3 represented 69.0% of the total, while R1 represented 31.0%, and the ratio of the amount of R3 consumed to the amount of R1 consumed was 2.2. The animals ate 41.5 grams/day/cat of R3, compared with 19.0 grams/day/cat of R1. Fifteen cats preferred R3 over R1, while two preferred R1 over R3, and one showed no preference.

Example 6

Eighteen adult cats were each offered a first test ration R2, and a second test ration R3 in a two bowl test over a period of two days. R2 was Meow Mix® basal formula kibbles coated with liquid animal digest at 3% by weight plus tetrasodium pyrophosphate at 0.5% by weight. R3 was Meow Mix® basal formula kibbles coated with liquid animal digest at 3% by weight plus

- 12 -

tetrasodium pyrophosphate at 0.75% by weight. Of all food consumed over the two days, R3 represented 60.9% of the total, while R2 represented 39.1%, and the ratio of the amount of R3 consumed to the amount of R2 consumed was 1.6. The animals ate 35.1 grams/day/cat of R3, compared with 23.2 grams/day/cat of R1. Fourteen cats preferred R3 over R2, while three preferred R2 over R3, and one showed no preference. The results suggest a positive dose response relationship between the amount of tetrasodium pyrophosphate added to the food and the degree of animal preference for the food.

Table 2 summarizes the results of Examples 5 and 6.

10

TABLE 2

Ex.	Ratio	most consumed / least consumed	% of all food consumed	Grams eaten /day/cat	# of animals with pref.	# of animals without pref.
5	R1	2.2	31.0	19.0	2	1
	R3		69.0	41.5	15	
6	R2	1.6	39.1	23.2	3	1
	R3		60.9	35.1	14	

Examples 1 through 6 show that tetrasodium pyrophosphate improves palatability in both dry and liquid formulations of a palatability enhancer. The addition of tetrasodium pyrophosphate to liquid animal digest was found to be more effective as a palatability enhancer than liquid animal digest alone. The addition of tetrasodium pyrophosphate to a dry cat coating produced a more effective palatability enhancer than a typical dry coating alone.

The use of tetrasodium pyrophosphate to enhance the palatability of dry and semi-dry pet foods represents a substantial improvement over current methods. The palatability enhancer is particularly effective for enhancing the palatability to cats of dry and semi-dry cat foods. Further, the use of tetrasodium pyrophosphate

- 13 -

represents a relatively low added production cost for pet food manufacturers. In addition, the improved palatability enhancing effect of tetrasodium pyrophosphate survives a range of different application systems and different formulations.

5 From the preceding description of various embodiments of the present invention, it is evident that the objects of the invention are attained. Although the invention has been described and illustrated in detail, it is to be clearly understood that the same is intended by way of illustration and example only and is not to be taken by way of limitation. Accordingly, the spirit and scope of the invention are to be limited only by the terms of the appended claims.

- 14 -

Claims:

1. A method for enhancing the palatability of an extruded pet food, said method comprising the steps of:
providing a palatability enhancing composition including tetrasodium
5 pyrophosphate; and
applying the palatability enhancer to the extruded pet food.
2. A method in accordance with Claim 1 wherein said step of applying the palatability enhancer comprises applying an amount of the palatability enhancer sufficient to apply about 0.1% to about 1.0% by weight tetrasodium pyrophosphate to the extruded pet food composition.
3. A method in accordance with Claim 1 wherein the palatability enhancing composition is dry and said step of applying the palatability enhancing composition to the extruded pet food comprises dusting the palatability enhancing composition onto the extruded pet food.
4. A method in accordance with Claim 1 wherein said step of applying the palatability enhancing composition comprises adding about 0.1% to about 3.0% by weight of the palatability enhancing composition to the extruded pet food.
5. A method in accordance with Claim 1 wherein the extruded pet food is a dry cat food.
6. A method in accordance with Claim 1 wherein the extruded pet food is a semi-dry cat food.

- 15 -

7. A method in accordance with Claim 1 wherein the palatability enhancing composition is a liquid and said step of applying the palatability enhancing composition to the extruded pet food comprises spraying the palatability enhancing composition onto the extruded pet food.

8. A method in accordance with Claim 1 wherein said step of providing a palatability enhancing composition comprises the steps of:

combining about 0.1% to about 99% by weight tetrasodium pyrophosphate, with dairy by-product, animal by-product, carbohydrate and microbial protein in a mixer; and

blending the tetrasodium pyrophosphate, dairy by-product, animal by-product, carbohydrate and microbial protein to homogeneity.

9. A palatability enhancing composition for extruded pet food, said composition comprising:

about 0.1% to about 99% by weight tetrasodium pyrophosphate;

about 20% to about 40% by weight microbial protein; and

about 20% to about 40% by weight animal by-product.

10. A palatability enhancing composition in accordance with Claim 9 further comprising about 15% to about 30% by weight carbohydrate.

11. A palatability enhancing composition in accordance with Claim 9 further comprising about 15% to about 30% by weight dairy by-product.

12. A palatability enhancing composition in accordance with Claim 9 comprising from about 5% to about 35% by weight tetrasodium pyrophosphate.

13. A palatability enhancing composition in accordance with Claim 9

- 16 -

comprising from about 18% to about 25% by weight tetrasodium pyrophosphate.

14. A palatability enhancing composition in accordance with Claim 9 wherein said palatability enhancing composition is dry.

15. A palatability enhancing composition in accordance with Claim 9 wherein said palatability enhancing composition is liquid.

16. A pet food comprising:
a nutritionally balanced extruded pet food composition; and
a palatability enhancing composition applied to a surface of said extruded
5 pet food composition, said palatability enhancing composition comprising from about 0.1% by weight to about 99% by weight tetrasodium pyrophosphate.

17. A pet food in accordance with Claim 16 wherein said extruded pet food composition comprises a dry cat food composition.

18. A pet food in accordance with Claim 16 wherein said extruded pet food composition comprises a semi-dry cat food composition.

19. A pet food in accordance with Claim 16 wherein said palatability enhancing composition includes about 18% to about 25% by weight tetrasodium pyrophosphate.

20. A pet food in accordance with Claim 16 wherein said palatability enhancing composition is dry.

21. A pet food in accordance with Claim 16 wherein said palatability enhancing composition is liquid.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/26640

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A21D 2/02; A23K 1/18

US CL : 426/549, 641, 656, 657, 658, 805

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 426/549, 641, 656, 657, 658, 805

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST/US PATENTS, STN/CAS MULTIFILE FOOD.

search terms: pyrophosphates, palatability, pet food, dog food, dog snacks, semi-moist.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,000,940 A (STAPLES et al.) 19 March 1991, see whole document	1-21
Y	JP 59001409 A (LION CORP.) 06 January 1984, see whole document	1-21
Y	US 5,186,964 A (GIERHART et al.) 16 February 1993, see whole document	1-21
Y	US 4,678,662 A (CHAN) 07 July 1987, see whole document	1-21

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

*	Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance		
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed	"&"	document member of the same patent family

Date of the actual completion of the international search

13 JANUARY 2000

Date of mailing of the international search report

03 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CHHAYA SAYALA

Telephone No. (703) 308-0661